

Serine/threonine-protein phosphatase 2A regulatory subunit B'


PP2A-2 (CAB036000)



Immunohistochemistry

IHC protocol: HIER pH 6, Dilution 1:1200

IHC test staining: Failed due to membranous and cytoplasmic staining in most tissues examined.



Western blot


WB Size markers (kDa): 230, 130, 95, 72, 56, 36, 28, 17, 11

WB Lanes: Marker(1), RT-4(2), U251 MG(3), Plasma(4), Liver(5), Tonsil(6)

WB Target weight (kDa): 37, 32, 30, 29, 28, 24, 19, 15, 14, 12, 7

WB Validation: Supportive (Band of predicted size in kDa (+/-20%) with additional bands present)


PP2A-3 (CAB036001)



Immunohistochemistry

IHC protocol: HIER pH 6, Dilution 1:1200

IHC test staining: Widely expressed slightly granular cytoplasmic staining. Strong staining was seen in e.g. testis, kidney, pancreas and fallopian tube. Some nuclear staining can also be seen.



Western blot

WB Size markers (kDa): 230, 130, 95, 72, 56, 36, 28, 17, 11

WB Lanes: Marker(1), RT-4(2), U251 MG(3), Plasma(4), Liver(5), Tonsil(6)

WB Target weight (kDa): 37, 32, 30, 29, 28, 24, 19, 15, 14, 12, 7

WB Validation: Supportive (Band of predicted size in kDa (+/-20%) with additional bands present)