CPTC-JUN-6 (CAB080220)

Uniprot ID: P05412

Protein name: JUN_HUMAN Full name: Transcription factor Jun


Tissue specificity: Expressed in the developing and adult prostate and prostate cancer cells. Function: Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:10995748, PubMed:22083952). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).

Subcellular location: Nucleus

Protein existence: Experimental evidence at protein level

Comment:

Immunohistochemistry




Immunofluorescence



IF Overlay:	antibody (green), anti-tubulin (red) and DAPI (blue)
IF main location:	Nucleoplasm - 3: Supportive (auto)
IF additional location:	Nuclear bodies - 3: Supportive (auto)
IF approved for publication on HPA:	Yes
IF in THP-1:	Nuclear bodies
IF in U-2 OS:	Nucleoplasm

Western blot



WB Size markers (kDa):	250, 130, 100, 70, 55, 35, 25, 15, 10	
WB Lanes:	Marker (1), RT4 (2), U-251 MG (3), Plasma (4), Liver (5), Tonsil (6)	
WB Target weight (kDa):	36	
WB Validation:	Uncertain (No bands detected.)	