

# CPTC-RB1-1 (CAB080142)

Uniprot ID: [P06400](#)

Protein name: RB\_HUMAN

Full name: Retinoblastoma-associated protein

Tissue specificity: Expressed in the retina. Expressed in foreskin keratinocytes (at protein level) (PubMed:20940255).

Function: Tumor suppressor that is a key regulator of the G1/S transition of the cell cycle (PubMed:10499802). The hypophosphorylated form binds transcription regulators of the E2F family, preventing transcription of E2F-responsive genes (PubMed:10499802). Both physically blocks E2Fs transactivating domain and recruits chromatin-modifying enzymes that actively repress transcription (PubMed:10499802). Cyclin and CDK-dependent phosphorylation of RB1 induces its dissociation from E2Fs, thereby activating transcription of E2F responsive genes and triggering entry into S phase (PubMed:10499802). RB1 also promotes the G0-G1 transition upon phosphorylation and activation by CDK3/cyclin-C (PubMed:15084261). Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1, KMT5B and KMT5C, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. Mediates transcriptional repression by SMARCA4/BRG1 by recruiting a histone deacetylase (HDAC) complex to the c-FOS promoter. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC1 repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex (By similarity). (Microbial infection) In case of viral infections, interactions with SV40 large T antigen, HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity.

Subcellular location:


Nucleus (*experimental evidence*)

NOTE: During keratinocyte differentiation, acetylation by KAT2B/PCAF is required for nuclear localization.

Protein existence: Experimental evidence at protein level


Comment:

## Immunohistochemistry




IHC protocol:	HIER pH6, Dilution 1:300
IHC test staining:	Weak cytoplasmic in a few tissues.
Literature conformance:	Not consistent with gene/protein characterization data
Literature significance:	
RNA similarity:	Very low consistency between antibody staining and RNA expression data
RNA tissue specificity:	Low tissue specificity
RNA tissue distribution:	Detected in all
IHC Sibling similarity:	Other antibody shows dissimilar IHC staining pattern

## Immunofluorescence



IF Overlay:	antibody (green), anti-tubulin (red) and DAPI (blue)
IF main location:	Cytosol - 12: <b>Uncertain</b> (auto) Nucleoplasm - 3: <b>Supportive</b> (auto)
IF additional location:	
IF approved for publication on HPA:	No
IF in SiHa:	Nucleoplasm Cytosol
IF in SK-MEL-30:	Nucleoplasm Cytosol
IF in U-2 OS:	Cytosol

# Western blot



<b>WB Size markers (kDa):</b>	250, 130, 100, 70, 55, 35, 25, 15, 10
<b>WB Lanes:</b>	Marker (1), RT4 (2), U-251 MG (3), Plasma (4), Liver (5), Tonsil (6)
<b>WB Target weight (kDa):</b>	11, 13, 105, 106
<b>WB Validation:</b>	Supported (Band of predicted size in kDa (+/-20%) with additional bands present.)