

CPTC-PP2A-2 (CAB080054)

Uniprot ID: [Q15257](#)

Protein name: PTPA_HUMAN

Full name: Serine/threonine-protein phosphatase 2A activator

Tissue specificity: Widely expressed.

Function: PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. Acts as a regulatory subunit for serine/threonine-protein phosphatase 2A (PP2A) modulating its activity or substrate specificity, probably by inducing a conformational change in the catalytic subunit, a proposed direct target of the PPlase. Can reactivate inactive phosphatase PP2A-phosphatase methylesterase complexes (PP2A(i)) in presence of ATP and Mg(2+) (By similarity). Reversibly stimulates the variable phosphotyrosyl phosphatase activity of PP2A core heterodimer PP2A(D) in presence of ATP and Mg(2+) (in vitro). The phosphotyrosyl phosphatase activity is dependent of an ATPase activity of the PP2A(D):PPP2R4 complex. Is involved in apoptosis; the function appears to be independent from PP2A.

Subcellular location:


Cytoplasm (*experimental evidence*)

Nucleus (*experimental evidence*)

Protein existence: Experimental evidence at protein level


Comment:

Immunohistochemistry




IHC protocol:	HIER pH6, Dilution 1:900
IHC test staining:	Mainly cytoplasmic positivity in most tissues.
Literature conformance:	Partly consistent with extensive gene/protein characterization data
Literature significance:	
RNA similarity:	Medium consistency between antibody staining and RNA expression data
RNA tissue specificity:	Low tissue specificity
RNA tissue distribution:	Detected in all
IHC Sibling similarity:	Other antibody shows similar IHC staining pattern

Immunofluorescence



IF Overlay:	antibody (green), anti-tubulin (red) and DAPI (blue)
IF main location:	Vesicles - 12: Uncertain (manual)
IF additional location:	
IF approved for publication on HPA:	No
IF in SiHa:	Vesicles
IF in SK-MEL-30:	Negative
IF in U-2 OS:	Negative

Western blot



WB Size markers (kDa):	250, 130, 100, 70, 55, 35, 25, 15, 10
WB Lanes:	Marker (1), RT4 (2), U-251 MG (3), Plasma (4), Liver (5), Tonsil (6)
WB Target weight (kDa):	5, 7, 7, 7, 7, 7, 7, 8, 12, 14, 14, 15, 16, 19, 24, 32, 33, 33, 33, 37, 37, 41, 42
WB Validation:	Supported (Band of predicted size in kDa (+/-20%) with additional bands present.)