

# CPTC-TP53-1 (CAB079964)

**Uniprot ID:** P04637

**Protein name:** P53\_HUMAN

**Full name:** Cellular tumor antigen p53

**Tissue specificity:** Ubiquitous. Isoforms are expressed in a wide range of normal tissues but in a tissue-dependent manner. Isoform 2 is expressed in most normal tissues but is not detected in brain, lung, prostate, muscle, fetal brain, spinal cord and fetal liver. Isoform 3 is expressed in most normal tissues but is not detected in lung, spleen, testis, fetal brain, spinal cord and fetal liver. Isoform 7 is expressed in most normal tissues but is not detected in prostate, uterus, skeletal muscle and breast. Isoform 8 is detected only in colon, bone marrow, testis, fetal brain and intestine. Isoform 9 is expressed in most normal tissues but is not detected in brain, heart, lung, fetal liver, salivary gland, breast or intestine.

**Function:** Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed:12524540). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed:12524540). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkn1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER2 (PubMed:24051492).

**Subcellular location:**

**Unnamed:**

Cytoplasm (*experimental evidence*)

Nucleus (*experimental evidence*)

Nucleus > PML body (*experimental evidence*)

Endoplasmic reticulum (*experimental evidence*)

Mitochondrion matrix (*experimental evidence*)

Cytoplasm > Cytoskeleton > Microtubule organizing center > Centrosome (*experimental evidence*)

**NOTE:** Interaction with BANP promotes nuclear localization (PubMed:15701641). Recruited into PML bodies together with CHEK2 (PubMed:12810724). Translocates to mitochondria upon oxidative stress (PubMed:22726440). Translocates to mitochondria in response to mitomycin C treatment (PubMed:27323408).

**Isoform 1:**

Nucleus

Cytoplasm

**NOTE:** Predominantly nuclear but localizes to the cytoplasm when expressed with isoform 4.

**Isoform 2:**

Nucleus

Cytoplasm

**NOTE:** Localized mainly in the nucleus with minor staining in the cytoplasm.

**Isoform 3:**

Nucleus

Cytoplasm

**NOTE:** Localized in the nucleus in most cells but found in the cytoplasm in some cells.

**Isoform 4:**

Nucleus

Cytoplasm

**NOTE:** Predominantly nuclear but translocates to the cytoplasm following cell stress.

**Isoform 7:**

Nucleus

Cytoplasm

**NOTE:** Localized mainly in the nucleus with minor staining in the cytoplasm.

**Isoform 8:**

Nucleus

Cytoplasm

**NOTE:** Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli.


**Isoform 9:**

Cytoplasm

**Protein existence:** Experimental evidence at protein level

**Comment:**


## Immunohistochemistry



<b>IHC protocol:</b>	HIER pH6, Dilution 1:250
<b>IHC test staining:</b>	Weak nuclear positivity in pancreas, remaining normal tissues were negative.
<b>Literature conformance:</b>	Not consistent with gene/protein characterization data
<b>Literature significance:</b>	
<b>RNA consistency:</b>	Not consistent with RNA expression data
<b>IHC Sibling similarity:</b>	Other antibody shows dissimilar IHC staining pattern
<b>IHC fail comment:</b>	ANTIBODY FAILED: Improbable histological location, Not consistent with RNA


## Immunofluorescence

<b>IF Overlay:</b>	antibody (green), anti-tubuline (red) and DAPI (blue)
<b>IF main location:</b>	



IF additional location:	
IF Antibody score:	Failed IF
IF in A549:	Negative
IF in HEK 293:	Negative
IF in U-2 OS:	Negative

# Western blot



<b>WB Size markers (kDa):</b>	250, 130, 100, 70, 55, 35, 25, 15, 10
<b>WB Lanes:</b>	Marker (1), RT4 (2), U-251 MG (3), Plasma (4), Liver (5), Tonsil (6)
<b>WB Target weight (kDa):</b>	3, 14, 15, 18, 18, 21, 21, 23, 24, 24, 27, 30, 32, 33, 34, 38, 38, 38, 39, 39, 39, 39, 43, 44, 44
<b>WB Validation:</b>	Uncertain (No bands detected.)